

*Multiple targets for flecainide action: implications for cardiac arrhythmogenesis*

Samantha C. Salvage<sup>1,2</sup>, Karthik H. Chandrasekharan<sup>2</sup>, Kamalan Jeevaratnam<sup>3,4</sup>, Angela F. Dulhunty<sup>5</sup>, Andrew J. Thompson<sup>6</sup>, Antony P. Jackson<sup>1</sup>, Christopher L-H Huang<sup>1,2</sup>

<sup>1</sup>Department of Biochemistry, Hopkins Building, University of Cambridge, Cambridge, CB2 1QW, United Kingdom.

<sup>2</sup>Physiological Laboratory, University of Cambridge, Downing Street, Cambridge, CB2 3EG, United Kingdom.

<sup>3</sup>Faculty of Health and Medical Science, VSM Building, University of Surrey, Guildford, GU2 7AL, United Kingdom.

<sup>4</sup>School of Medicine, Perdana University – Royal College of Surgeons Ireland, 43400 Serdang, Selangor Darul Ehsan, Malaysia.

<sup>5</sup>Muscle Research Group, Division of Molecular Bioscience, John Curtin School of Medical Research, Australian National University, Acton ACT 2601, Australia.

<sup>6</sup>Department of Pharmacology, University of Cambridge, Cambridge CB2 1PD, United Kingdom.

Address for correspondence: Christopher L-H Huang, Physiological Laboratory, University of Cambridge, Downing Street, Cambridge, CB2 3EG, United Kingdom. Phone 01223 333822, email: [clh11@cam.ac.uk](mailto:clh11@cam.ac.uk).

Keywords: flecainide, propafenone, tetracaine, lignocaine, mexiletine, Nav channel, Ryanodine receptor, cardiac arrhythmia, conduction velocity, triggering events.

Short title: Cardiac actions of flecainide